(19) World Intellectual Property Organization International Bureau



- I BORN BUNKKU KARUM KUMI BUNK BUNK BIRK I DIN OTOK BUNK BUNK BUNK BURK BUNK BUNK BUNK BURK BURK BURK BURK BU

(43) International Publication Date 15 January 2004 (15.01.2004)

PCT

(10) International Publication Number WO 2004/004734 A1

- (51) International Patent Classification⁷: A61K 31/519, A61P 1/00
- (21) International Application Number:

PCT/GB2003/002974

- (22) International Filing Date: 9 July 2003 (09.07.2003)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0216027.3 10 July 2002 (10.07.2002) GB 0304648.9 28 February 2003 (28.02.2003) GB

- (71) Applicant (for all designated States except US): ARACH-NOVA THERAPEUTICS LTD. [GB/GB]; 95 Halkett Place, St. Helier, Jersey JE1 1BX (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): CAVALLA, David [GB/GB]; Arachnova Ltd., St. John's Innovation Centre, Cambridge CB4 0WS (GB). GRISTWOOD, Robert, William [GB/GB]; Arachnova Ltd., St. John's Innovation Centre, Cambridge CB4 0WS (GB).

- (74) Agent: GILL, Jennings & Every; Broadgate House, 7 Elton Street, London EC2M 7LH (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 4-(2-FLUOROPHENYL)-6-METHYL-2(1-PIPERAZINYL)THIENO(2,3-D) PYRIMIDINE IN THE TREATMENT OF FUNCTIONAL BOWEL DISORDER

(57) Abstract: Use of (4-(2-fluorophenyl)-6-methyl-2-(1-piperazinyl)thieno[2,3-D]pyrimidine or a salt thereof for the manufacture of a medicament for the treatment of a functional bowel disorder.

AL2

2004/004734 A1

10

15

20

25

35

4-(2-FLUOROPHENYL)-6-METHYL-2-(1-PIPERAZINYL)THIENO(2,3-D)PYRIMIDINE IN THE TREATMENT OF FUNCTIONAL BOWEL DISORDER

Field of the Invention

This invention relates to a new use for a known compound.

5 Background of the Invention

4-(2-Fluorophenyl)-6-methyl-2-(1-piperazinyl)thieno[2,3-D]pyrimidine monohydrate hydrochloride is known (see US-A-4695568) and has shown activity as an antidepressant. It has serotonin and noradrenergic reuptake blocking properties and this is thought to be the mechanism for its action as an antidepressant. The compound also has 5HT-3 blocking activity.

Functional bowel disorders are very common and include irritable bowel syndrome (IBS) and functional dyspepsia. IBS is the most common disorder diagnosed by gastroenterologists and one of the more common encountered in general practice. The overall prevalence rate is similar (approx 10%) in most industrialised countries. Some estimates of prevalence have reached 20%. The illness has a large economic impact on health care use and indirect costs, chiefly through absenteeism.

IBS falls into two categories of equal prevalence, constipation-predominant and diarrhoea-predominant. The available treatments are generally poor.

A recent approach to treating diarrhoea-predominant IBS has involved the use of alosetron. This drug works by blocking the 5HT-3 receptor. Other drugs with this mechanism of action have shown some limited activity in this disease, including granisetron. Alosetron, although effective, was withdrawn due to side-effects on the colon.

A recent approach to treating constipation-predominant IBS involved agonising the 5HT4 receptor. Two such agonists are in clinical trials, i.e. tegaserod and prucalopride. Other approaches being explored include using 5HT1 agonists such as buspirone.

Functional dyspepsia is characterised by impaired accommodation of the stomach to a meal and epigastric pain discomfort or pain. There is often early satiety and weight loss. The disorder is not well understood. Treatments include antispasmodics and drugs affecting gut motility. Early studies suggest that buspirone and serotonin reuptake inhibitors may be useful.

Summary of the Invention

Surprisingly, it has been found that the known compound identified above (referred to herein as MCI-225) has activity in the treatment of functional bowel

5

10

15

20

25

30

35

disorders. Its combination of serotonin and noradrenergic reuptake blockade and 5HT-3 receptor blockade has not previously been clearly identified as being responsible for activity in functional bowel disorders. Furthermore MCI-225, at doses effective in the treatment of bowel disorders, can produce a lower incidence of some of the side-effects which are commonly known to be associated with the clinical use of selective serotonin reuptake inhibitors, for example the production of nausea and vomiting or the induction of sexual dysfunction. It will be appreciated that any suitable form of the active principle may be used, e.g. another salt form, or a prodrug or active metabolite.

Description of Preferred Embodiments

By means of this invention, functional bowel disorders and associated pain symptoms can be treated, e.g. controlled or prevented. Such disorders include irritable bowel syndrome, including diarrhoea-predominant, constipation-predominant, and alternating constipation/diarrhoea IBS. The patient may be male or female, diarrhoea-

predominant IBS being particularly associated with women.

For use in the invention, the active compound can be formulated in any suitable manner together with a conventional diluent or carrier. The active compound is preferably administered by the oral route; other suitable routes of administration include sublingual/buccal, transdermal, intramuscular, intranasal, rectal, parenteral, subcutaneous, pulmonary and topical. An effective dose of the active agent will depend on the nature and degree of the complaint, the age and condition of the patient and other factors known to those skilled in the art. A typical daily dosage may be 0.1 mg to 1 g.

A pharmaceutical composition containing the active ingredient may be in the form of a sublingual tablet or patch. Suitable compositions for oral use include tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, syrups and elixirs. Suitable additives include sweetening agents, flavouring agents, colouring agents and preserving agents. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients, e.g. inert diluents such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as

5

10

15

20

25

30

35

glyceryl monostearate or glyceryl distearate may be employed. They may also be coated, to form osmotic therapeutic tablets for controlled release. Hard gelatin capsules may include an inert solid diluent, for example calcium carbonate, calcium phosphate or kaolin; soft gelatin capsules may include water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

The data on which this invention is based will now be described. In a study using intact animals, the ability of a drug to inhibit the reflex depressor response to colorectal distension can be assessed. In this model, an inhibition of the reflex indicates modulation of visceral nociceptive neurotransmission and, therefore, the use of the drug in functional bowel disease (e.g. IBS); see Kozlowski *et al*, 2000, Gut 46, 474-480. Allodynia and visceral pain are important components of functional bowel disease.

Study

Experiments were performed on male Sprague-Dawley rats (250-300 g). Anaesthesia was induced with isoflurane (2.5% in oxygen) and maintained with alpha chlorolose (80 mg/kg i.v.). The left carotid artery was cannulated for the measurement of blood pressure and heart rate and the left jugular vein cannulated for drug administration. A tracheal cannula was implanted for artificial respiration if required. A 10 mm long latex balloon was inserted intrarectally so that the tip of the balloon was 20 mm from the anal verge (Kozlowski *et al*, *supra*). The balloon was connected via a double lumen cannula to a pressure transducer and also to a saline-filled syringe for inflation/deflation of the balloon. Throughout the experiment, body temperature was kept constant at 36-38 C using a homeothermic blanket.

Once stable baseline parameters were obtained (approximately after 20 minutes), the balloon was rapidly inflated with increasing volumes of saline (0.5-2.5 ml) for 30 seconds at 5 minute intervals, and the resultant change in blood pressure recorded. Three distinct response curves were constructed, with a 10 minute stabilisation period between each curve. In one group of animals, 10 minutes prior to the commencement of the final distension response curve, a single bolus of MCI-225 (3 mg/kg) was administered intravenously; in a second group of animals, a single bolus dose of vehicle was administered. The effect of MCI-225 and vehicle was determined by analysing the changes in colorectal distension that evoked depressor response.

Falls in arterial blood pressure (mean absolute decreases in mean arterial pressure in mmHg, with standard error of mean in brackets) evoked by distension of the balloon, before adding drug, at 0.5, 1.0, 1.5, 2.0 and 2.5 ml balloon volume were 2.7

WO 2004/004734 PCT/GB2003/002974

(1.9), 12.4 (5.9), 24.0 (8.9), 36.3 (4.8) and 43.4 (6.0), respectively (all except final value n=6, final value n=5). Following administration of MCI-225 at 3 mg/kg i.v., the corresponding values were 2.2 (1.65), 6.3 (2.6), 10.6 (3.9), 15.3 (5.4) and 24.6 (7.3), respectively (all values except final value n=6, final value n=5).

The results clearly show that MCI-225 inhibited the distension-induced falls in blood pressure. The falls in blood pressure evoked by 2.0 and 2.5 ml balloon volumes were reduced with statistical significance following administration of MCI-225 at 3mg/kg, with p values (paired t test) of less than 0.01 and less than 0.05 respectively.

5

CLAIMS

- 1. Use of (4-(2-fluorophenyl)-6-methyl-2-(1-piperazinyl)thieno[2,3-D]pyrimidine or a salt thereof for the manufacture of a medicament for the treatment of a functional bowel disorder.
- 5 2. Use according to claim 1, wherein the salt is the hydrochloride monohydrate.
 - 3. Use according to claim 1 or claim 2, wherein the disorder is irritable bowel syndrome.
 - 4. Use according to claim 3, wherein the disorder is diarrhoea-predominant irritable bowel syndrome.
- 10 5. Use according to claim 4, wherein the disorder is in a female patient.
 - 6. Use according to claim 3, wherein the disorder is alternating constipation/diarrhoea irritable bowel syndrome.
 - 7. Use according to claim 3, wherein the disorder is constipation-predominant irritable bowel syndrome.

INTERNATIONAL SEARCH REPORT

PCT/GB 03/02974

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 A61K31/519 A61P1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, EMBASE, BIOSIS, CHEM ABS Data

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EGUCHI J ET AL: "The anxiolytic-like effect of MCI-225, a selective NA reuptake inhibitor with 5-HT3 receptor antagonism" PHARMACOLOGY BIOCHEMISTRY AND BEHAVIOR, ELSEVIER, US, vol. 68, no. 4, April 2001 (2001-04), pages 677-683, XP002239887 ISSN: 0091-3057 abstract	1-7

X Patent family members are listed in annex.
"T" later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention. "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone. "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family
Date of mailing of the International search report
. 17/10/2003
Authorized officer Sk.jöldebrand, C

1

INTERNATIONAL SEARCH REPORT

PCT/GB 03/02974

	ation) DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with Indication, where appropriate, of the relevant passages	Relevant to claim No.		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Tielevalit to Clauti No.		
1	CROWELL M D: "The role of serotonin in the pathophysiology of irritable bowel syndrome" AMERICAN JOURNAL OF MANAGED CARE 2001 UNITED STATES, vol. 7, no. 8 SUPPL., 2001, pages S252-S260, XP009018426 ISSN: 1088-0224 the whole document	1-7		
ſ	US 6 284 770 B1 (MANGEL ALLEN WAYNE ET AL) 4 September 2001 (2001-09-04) the whole document	1-7		
4	DE 100 63 223 A (MERCK PATENT GMBH) 20 June 2002 (2002-06-20) abstract page 2, line 43	1-7		
4	US 4 695 568 A (NINOMIYA KUNIHIRO ET AL) 22 September 1987 (1987-09-22) cited in the application the whole document	1-7		
•				
	•			

INTERNATIONAL SEARCH REPORT

Information on patent family members

PCT/GB 03/02974

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
US 6284770	B1	04-09-2001	AU	750818		25-07-2002
			ΑU	9629398	Α	27-04-1999
			BR	9812886	Α	08-08-2000
			CA	2305751	A1	15-04-1999
			EE	200000214	Α	15-06-2001
			EP	1021174	A2	26-07-2000
			HR	20000198	A1	30-04-2001
			HU	0003750	A2	28-10-2001
			JP	2001518495	T	16-10-2001
			NO	20001776	Α	06-06-2000
			NZ	503698	Α	25-10-2002
			PL	340337	A1	29-01-2001
			SK	4862000	A3	18-01-2001
			CN	1281357	T	24-01-2001
			WO	9917755	A2	15-04-1999
			TR	200000913	T2	22-01-2001
			US	2003036549	A1	20-02-2003
			บร	2001044450		22-11-2001
			ZA	9809061		05-07-2001
DE 10063223	Α	20-06-2002	DE	10063223	A1	20-06-2002
			ΑU	2795702	Α	01-07-2002
			CA	2431074	A1	27-06-2002
			CZ	20031754	A3	17-09-2003
			WO	0249650	.A2	27-06-2002
			EP	1347761	A2	01-10-2003
			NO	20032772	A	18-06-2003
US 4695568	. A	22-09-1987	JP	1699365	C	28-09-1992
	•		JP	3067071		21-10-1991
			JP	60146891		02-08-1985
			ΑT		Τ.	15-07-1988
			CA	1224782		28-07-1987
_			DE	3472106		21-07-1988
•			DK		А ,В,	06-07-1985
			EP	0150469	A1	07-08-1985
			HU	37435	A2	28-12-1985